

A Simple Construction of a Triangle from its Centroid, Incenter, and a Vertex

Eric Danneels

Abstract. We give a simple ruler and compass construction of a triangle given its centroid, incenter, and one vertex. An analysis of the number of solutions is also given.

1. Construction

The ruler and compass construction of a triangle from its centroid, incenter, and one vertex was one of the unresolved cases in [3]. An analysis of this problem, including the number of solutions, was given in [1]. In this note we give a very simple construction of triangle ABC with given centroid G , incenter I , and vertex A . The construction depends on the following propositions. For another slightly different construction, see [2].

Proposition 1. *Given triangle ABC with Nagel point N , let D be the midpoint of BC . The lines ID and AN are parallel.*

Proof. The centroid G divides each of the segments AD and NI in the ratio $AG : GD = NG : GI = 2 : 1$. See Figure 1. □




Figure 1




Figure 2

Proposition 2. *Let X be the point of tangency of the incircle with BC . The antipode of X on the circle with diameter ID is a point on AN .*

Publication Date: April 12, 2005. Communicating Editor: Paul Yiu.

The author wishes to thank Paul Yiu for his help in the preparation of this paper.

Proof. This follows from the fact that the antipode of X on the incircle lies on the segment AN . See Figure 2. \square

Construction. Given G , I , and A , extend AG to D such that $AG : GD = 2 : 1$. Construct the circle \mathcal{C} with diameter ID , and the line \mathcal{L} through A parallel parallel to ID .

Let Y be an intersection of the circle \mathcal{C} and the line \mathcal{L} , and X the antipode of Y on \mathcal{C} such that A is outside the circle $I(X)$. Construct the tangents from A to the circle $I(X)$. Their intersections with the line DX at the remaining vertices B and C of the required triangle. See Figure 3.




Figure 3

2. Number of solutions

We set up a Cartesian coordinate system such that $A = (0, 2k)$ and $I = (0, -k)$. If $G = (u, v)$, then $D = \frac{1}{2}(3G - A) = (\frac{3}{2}u, \frac{3}{2}v - k)$. The circle \mathcal{C} with diameter ID has equation

$$2(x^2 + y^2) - 3ux - (3v - 4k)y + (2k^2 - 3kv) = 0$$

and the line \mathcal{L} through A parallel to ID has slope $\frac{v}{u}$ and equation

$$vx - uy + 2ku = 0.$$

The line \mathcal{L} and the circle \mathcal{C} intersect at 0, 1, 2 real points according as

$$\Delta := (u^2 + v^2 - 4ku)(u^2 + v^2 + 4ku)$$

is negative, zero, or positive. Since $x^2 + y^2 \pm 4kx = 0$ represent the two circles of radii $2k$ tangent to each other externally and to the y -axis at $(0, 0)$, Δ is negative,

zero, or positive according as G lies in the interior, on the boundary, or in the exterior of the union of the two circles.

The intersections of the circle and the line are the points

$$Y_\varepsilon = \left(\frac{3u(u^2 + v^2 - 4kv - \varepsilon\sqrt{\Delta})}{4(u^2 + v^2)}, \frac{8k(u^2 + v^2) + 3v(u^2 + v^2 - 4kv - \varepsilon\sqrt{\Delta})}{4(u^2 + v^2)} \right)$$

for $\varepsilon = \pm 1$. Their antipodes on \mathcal{C} are the points

$$X_\varepsilon = \left(\frac{3u(u^2 + v^2 + 4kv + \varepsilon\sqrt{\Delta})}{4(u^2 + v^2)}, \frac{-16k(u^2 + v^2) + 3v(u^2 + v^2 + 4kv + \varepsilon\sqrt{\Delta})}{4(u^2 + v^2)} \right).$$

There is a triangle ABC tritangent to the circle $I(X_\varepsilon)$ and with DX_ε as a side-line if and only if the point A lies outside the circle $I(X_\varepsilon)$. Note that $IA = 3k$ and

$$IX_+^2 = \frac{9}{8}(u^2 + v^2 + \sqrt{\Delta}), \quad IX_-^2 = \frac{9}{8}(u^2 + v^2 - \sqrt{\Delta}).$$

From these, we make the following conclusions.

- (i) If $u^2 + v^2 - 8k^2 \geq \sqrt{\Delta}$, then A lies inside or on $I(X_-)$. In this case, there is no triangle.
- (ii) If $-\sqrt{\Delta} \leq u^2 + v^2 - 8k^2 < \sqrt{\Delta}$, then A lies outside $I(X_-)$ but not $I(X_+)$. There is exactly one triangle.
- (iii) If $u^2 + v^2 - 8k^2 < -\sqrt{\Delta}$, then A lies outside $I(X_+)$ (and also $I(X_-)$). There are in general two triangles.

It is easy to see that the condition $-\sqrt{\Delta} < u^2 + v^2 - 8k^2 < \sqrt{\Delta}$ is equivalent to $(v - 2k)(v + 2k) > 0$, i.e., $|v| > 2k$. We also note the following.

- (i) When the line D_ε passes through A , the corresponding triangle degenerates. The condition for collinearity leads to

$$u(3u^2 + 3v^2 - 4kv \pm \sqrt{\Delta}) = 0.$$

Clearly, $u = 0$ gives the y -axis. The corresponding triangle is isosceles. On the other hand, the condition $3u^2 + 3v^2 - 4kv \pm \sqrt{\Delta} = 0$ leads to

$$(u^2 + v^2)(u^2 + v^2 - 3kv + 2k^2) = 0,$$

i.e., (u, v) lying on the circle tangent to the circles $x^2 + y^2 \pm 4kx = 0$ at $(\pm \frac{2k}{5}, \frac{6k}{5})$ and the line $y = 2k$ at A .

- (ii) If $v > 0$, the circle $I(X_\varepsilon)$, instead of being the incircle, is an excircle of the triangle. If G lies inside the region $ATOT'A$ bounded by the circular segments, one of the excircles is the A -excircle. Outside this region, the excircle is always a B/C -excircle.

From these we obtain the distribution of the position of G , summarized in Table 1 and depicted in Figure 4, for the various numbers of solutions of the construction problem. In Figure 4, the number of triangles is

- 0 if G in an unshaded region, on a dotted line, or at a solid point other than I ,
- 1 if G is in a yellow region or on a solid red line,
- 2 if G is in a green region.

Table 1. Number N of non-degenerate triangles according to the location of G relative to A and I

N	Location of centroid $G(u, v)$
0	$(0, 0), (\pm 2k, 2k);$ $(\pm \frac{2k}{5}, \frac{6k}{5});$ $v = 2k;$ $ u > 2k - \sqrt{4k^2 - v^2}, -2k \leq v < 2k.$
1	$u = 0, 0 < v < 2k;$ $-2k < u < 2k, v = -2k;$ $u = 2k - \sqrt{4k^2 - v^2}, 0 < v < 2k;$ $ v > 2k;$ $u^2 + v^2 - 3kv + 2k^2 = 0$ except $(0, 2k), (\pm \frac{2k}{5}, \frac{6k}{5}).$
2	$ u < 2k - \sqrt{4k^2 - v^2}, 0 < v < 2k,$ but $u^2 + v^2 - 3kv + 2k^2 \neq 0.$




Figure 4

References

- [1] J. Anglesio and V. Schindler, Problem 10719, *Amer. Math. Monthly*, 106 (1999) 264; solution, 107 (2000) 952–954.
- [2] E. Danneels, Hyacinthos message 11103, March 22, 2005.
- [3] W. Wernick, Triangle constructions with three located points, *Math. Mag.*, 55 (1982) 227–230.

Eric Danneels: Hubert d'Ydewallestraat 26, 8730 Beernem, Belgium
E-mail address: eric.danneels@pandora.be